

Material

- ▶ Glad & Ljung Ch. 12.2
- ▶ Khalil Ch. 4.1-4.3
- ▶ Slotine and Li: Chapter 3 (not 3.5.2-3.5.3)
- ▶ Lecture notes

Alexandr Mihailovich Lyapunov (1857–1918)



Master thesis “On the stability of ellipsoidal forms of equilibrium of rotating fluids,” St. Petersburg University, 1884.

Doctoral thesis “The general problem of the stability of motion,” 1892.

Examples

Start with a Lyapunov *candidate* V to measure e.g.,

- ▶ “size”¹ of state and/or output error,
- ▶ “size” of deviation from true parameters,
- ▶ energy difference from desired equilibrium,
- ▶ weighted combination of above
- ▶ ...


Example of common choice in adaptive control

$$V = \frac{1}{2} (e^2 + \gamma_a \tilde{a}^2 + \gamma_b \tilde{b}^2)$$

(here weighted sum of output error and parameter errors)

¹ Often a magnitude measure or (squared) norm like $\|e\|_2, \dots$

A Motivating Example



$$m\ddot{x} = - \underbrace{b\dot{x}}_{\text{damping}} - \underbrace{k_0x - k_1x^3}_{\text{spring}}$$

$b, k_0, k_1 > 0$

Total energy = kinetic + pot. energy: $V = \frac{m\dot{x}^2}{2} + \int_0^x F_{\text{spring}} ds \Rightarrow$

$$V(x, \dot{x}) = m\dot{x}^2/2 + k_0x^2/2 + k_1x^4/4 > 0, \quad V(0, 0) = 0$$

$$\begin{aligned} \frac{d}{dt} V(x, \dot{x}) &= m\dot{x}\ddot{x} + k_0x\dot{x} + k_1x^3\dot{x} = \{\text{plug in system dynamics}^2\} \\ &= -b|\dot{x}|^3 < 0, \text{ for } \dot{x} \neq 0 \end{aligned}$$

What does this mean?

² Also referred to evaluate “along system trajectories”.

To be able to

- ▶ prove local and global stability of an equilibrium point using Lyapunov’s method
- ▶ show stability of a set (e.g., an equilibrium, or a limit cycle) using La Salle’s invariant set theorem.

Main idea

Lyapunov formalized the idea:

If the total energy is dissipated, then the system must be stable.

Main benefit: By looking at **how** an energy-like function V (a so called *Lyapunov function*) **changes over time**, we might **conclude** that a system is stable or asymptotically stable **without solving** the nonlinear differential equation.

Main question: How to find a Lyapunov function?

Analysis: Check if V is decreasing with time

- ▶ Continuous time: $\frac{dV}{dt} < 0$
- ▶ Discrete time: $V(k+1) - V(k) < 0$

Synthesis: Choose e.g. control law and/or parameter update law to satisfy $\dot{V} \leq 0$

$$\begin{aligned} \frac{dV}{dt} &= e\dot{e} + \gamma_a \tilde{a}\dot{\tilde{a}} + \gamma_b \tilde{b}\dot{\tilde{b}} = \\ &= \tilde{x}(-a\tilde{x} - \tilde{a}\tilde{x} + \tilde{b}u) + \gamma_a \tilde{a}\dot{\tilde{a}} + \gamma_b \tilde{b}\dot{\tilde{b}} = \dots \end{aligned}$$

If a is constant and $\tilde{a} = a - \hat{a}$ then $\dot{\tilde{a}} = -\dot{\hat{a}}$.

Choose update law $\frac{d\hat{a}}{dt}$ in a “good way” to influence $\frac{dV}{dt}$.
(more on this later...)

Stability Definitions

An equilibrium point $x = 0$ of $\dot{x} = f(x)$ is

- ▶ **locally stable**, if for every $R > 0$ there exists $r > 0$, such that

$$\|x(0)\| < r \Rightarrow \|x(t)\| < R, \quad t \geq 0$$

- ▶ **locally asymptotically stable**, if locally stable and

$$\|x(0)\| < r \Rightarrow \lim_{t \rightarrow \infty} x(t) = 0$$

- ▶ **globally asymptotically stable**, if asymptotically stable for all $x(0) \in \mathbf{R}^n$.

Lyapunov Theorem for Local Stability

Theorem Let $\dot{x} = f(x)$, $f(0) = 0$ where $x = 0$ is in the interior of $\Omega \subset \mathbb{R}^n$. Assume that $V : \Omega \rightarrow \mathbb{R}$ is a C^1 function. If

- (1) $V(0) = 0$
 - (2) $V(x) > 0$, for all $x \in \Omega$, $x \neq 0$
 - (3) $\dot{V}(x) \leq 0$ along all trajectories of the system in Ω
- $\Rightarrow x = 0$ is locally stable.

Furthermore, if also

- (4) $\dot{V}(x) < 0$ for all $x \in \Omega$, $x \neq 0$
- $\Rightarrow x = 0$ is locally asymptotically stable.

Conservation and Dissipation

Conservation of energy: $\dot{V}(x) = \frac{\partial V}{\partial x} f(x) = 0$, i.e., the vector field $f(x)$ is everywhere orthogonal to the normal $\frac{\partial V}{\partial x}$ to the level surface $V(x) = c$.

Example: Total energy of a lossless mechanical system or total fluid in a closed system.

Dissipation of energy: $\dot{V}(x) = \frac{\partial V}{\partial x} f(x) \leq 0$, i.e., the vector field $f(x)$ and the normal $\frac{\partial V}{\partial x}$ to the level surface $\{z : V(z) = c\}$ make an obtuse angle (Sw. "trubbig vinkel").

Example: Total energy of a mechanical system with damping or total fluid in a system that leaks.

Boundedness:

For any trajectory $x(t)$

$$V(x(t)) = V(x(0)) + \int_0^t \dot{V}(x(\tau)) d\tau \leq V(x(0))$$

which means that the whole trajectory lies in the set

$$\{z \mid V(z) \leq V(x(0))\}$$

For stability it is thus important that the sublevel sets $\{z \mid V(z) \leq c\}$ bounded $\forall c \geq 0 \iff V(x) \rightarrow \infty$ as $\|x\| \rightarrow \infty$.

Positive Definite Matrices

Definition: Symmetric matrix $M = M^T$ is

- ▶ **positive definite** ($M > 0$) if $x^T M x > 0$, $\forall x \neq 0$
- ▶ **positive semidefinite** ($M \geq 0$) if $x^T M x \geq 0$, $\forall x$

Lemma:

- ▶ $M = M^T > 0 \iff \lambda_i(M) > 0$, $\forall i$
- ▶ $M = M^T \geq 0 \iff \lambda_i(M) \geq 0$, $\forall i$

$$M = M^T > 0 \quad V(x) := x^T M x$$

\Downarrow

$$V(0) = 0, \quad V(x) > 0, \quad \forall x \neq 0$$

$V(x)$ candidate Lyapunov function


Lyapunov Functions (\approx Energy Functions)

A function V that fulfills (1)–(3) is called a *Lyapunov function*.


Condition (3) means that V is non-increasing along all trajectories in Ω :

$$\dot{V}(x) = \frac{\partial V}{\partial x} \dot{x} = \sum_i \frac{\partial V}{\partial x_i} f_i(x) \leq 0$$

$$\text{where } \frac{\partial V}{\partial x} = \left[\frac{\partial V}{\partial x_1}, \frac{\partial V}{\partial x_2}, \dots, \frac{\partial V}{\partial x_n} \right]$$



Geometric interpretation



Vector field points into sublevel sets

Trajectories can only go to lower values of $V(x)$


2 min exercise—Pendulum

Show that the origin is locally stable for a mathematical pendulum.

$$\dot{x}_1 = x_2, \quad \dot{x}_2 = -\frac{g}{\ell} \sin x_1$$

Use as a Lyapunov function candidate

$$V(x) = (1 - \cos x_1)g\ell + \ell^2 x_2^2 / 2$$



More matrix results

- ▶ for symmetric matrix $M = M^T$

$$\lambda_{\min}(M) \|x\|^2 \leq x^T M x \leq \lambda_{\max}(M) \|x\|^2, \quad \forall x$$

Proof idea: factorize $M = U \Lambda U^T$, unitary U (i.e., $\|Ux\| = \|x\| \forall x$), $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$

- ▶ for any matrix M

$$\|Mx\| \leq \sqrt{\lambda_{\max}(M^T M)} \|x\|, \quad \forall x$$

Example- Lyapunov function for linear system


$$\dot{x} = Ax = \begin{bmatrix} -1 & 4 \\ 0 & -3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \quad (1)$$

Eigenvalues of A : $\{-1, -3\} \Rightarrow$ (global) asymptotic stability.

Find a quadratic Lyapunov function for the system (??):

$$V(x) = x^T P x = \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \quad P = P^T > 0$$

Take any $Q = Q^T > 0$, say $Q = I_{2 \times 2}$. Solve $A^T P + PA = -Q$.



Phase plot showing that

$$V = \frac{1}{2}(x_1^2 + x_2^2) = \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 0.5 & 0 \\ 0 & 0.5 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \text{ does NOT work.}$$

Lyapunov Stability for Linear Systems

Linear system: $\dot{x} = Ax$

Lyapunov equation: Let $Q = Q^T > 0$. Solve

$$PA + A^T P = -Q$$

with respect to the symmetric matrix P .

Lyapunov function: $V(x) = x^T P x, \Rightarrow$

$$\dot{V}(x) = x^T P \dot{x} + \dot{x}^T P x = x^T (PA + A^T P) x = -x^T Q x < 0$$

Asymptotic Stability: If $P = P^T > 0$, then the Lyapunov Stability Theorem implies (local=global) asymptotic stability, hence the eigenvalues of A must satisfy $\text{Re } \lambda_k(A) < 0, \forall k$

Interpretation

Assume $\dot{x} = Ax, x(0) = z$. Then

$$\int_0^\infty x^T(t) Q x(t) dt = z^T \left(\int_0^\infty e^{A^T t} Q e^{At} dt \right) z = z^T P z$$

Thus $V(z) = z^T P z$ is the cost-to-go from z (with no input) and integral quadratic cost function with weighting matrix Q .

Example cont'd

$$A^T P + PA = -I$$

$$\begin{bmatrix} -1 & 0 \\ 4 & -3 \end{bmatrix} \begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix} + \begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix} \begin{bmatrix} -1 & 4 \\ 0 & -3 \end{bmatrix} = \begin{bmatrix} -2p_{11} & -4p_{12} + 4p_{11} \\ -4p_{12} + 4p_{11} & 8p_{12} - 6p_{22} \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} \quad (2)$$


Solving for p_{11}, p_{12} and p_{22} gives

$$2p_{11} = -1$$

$$-4p_{12} + 4p_{11} = 0$$

$$8p_{12} - 6p_{22} = -1$$

$$\Rightarrow \begin{bmatrix} p_{11} & p_{12} \\ p_{12} & p_{22} \end{bmatrix} = \begin{bmatrix} 1/2 & 1/2 \\ 1/2 & 5/6 \end{bmatrix} > 0$$



Phase plot with level curves $x^T P x = c$ for P found in example.

Converse Theorem for Linear Systems

If $\text{Re } \lambda_k(A) < 0 \forall k$, then for every $Q = Q^T > 0$ there exists $P = P^T > 0$ such that $PA + A^T P = -Q$

Proof: Choose $P = \int_0^\infty e^{A^T t} Q e^{At} dt$. Then

$$\begin{aligned} A^T P + PA &= \lim_{t \rightarrow \infty} \int_0^t (A^T e^{A^T \tau} Q e^{A\tau} + e^{A^T \tau} Q A e^{A\tau}) d\tau \\ &= \lim_{t \rightarrow \infty} [e^{A^T \tau} Q e^{A\tau}]_0^t \\ &= -Q \end{aligned}$$

Lyapunov's Linearization Method

Recall from Lecture 2:

Theorem Consider

$$\dot{x} = f(x)$$

Assume that $f(0) = 0$. Linearization

$$\dot{x} = Ax + g(x), \quad \|g(x)\| = o(\|x\|) \text{ as } x \rightarrow 0.$$

(1) $\text{Re } \lambda_k(A) < 0, \forall k \Rightarrow x = 0$ locally asympt. stable

(2) $\exists k : \text{Re } \lambda_k(A) > 0 \Rightarrow x = 0$ unstable

Proof of (1) in Lyapunov's Linearization Method

Put $V(x) := x^T P x$. Then, $V(0) = 0$, $V(x) > 0 \forall x \neq 0$, and

$$\begin{aligned}\dot{V}(x) &= x^T P f(x) + f^T(x) P x \\ &= x^T P [Ax + g(x)] + [x^T A^T + g^T(x)] P x \\ &= x^T (PA + A^T P)x + 2x^T P g(x) = -x^T Q x + 2x^T P g(x)\end{aligned}$$

$$x^T Q x \geq \lambda_{\min}(Q) \|x\|^2$$

and for all $\gamma > 0$ there exists $r > 0$ such that

$$\|g(x)\| < \gamma \|x\|, \quad \forall \|x\| < r$$

Thus, choosing γ sufficiently small gives


$$\dot{V}(x) \leq -(\lambda_{\min}(Q) - 2\gamma \lambda_{\max}(P)) \|x\|^2 < 0$$

Radial Unboundedness is Necessary

If the condition $V(x) \rightarrow \infty$ as $\|x\| \rightarrow \infty$ is not fulfilled, then global stability cannot be guaranteed.

Example Assume $V(x) = x_1^2/(1+x_1^2) + x_2^2$ is a Lyapunov function for a system. Can have $\|x\| \rightarrow \infty$ even if $\dot{V}(x) < 0$.

Contour plot $V(x) = C$:



Example [Khalil]:

$$\begin{aligned}\dot{x}_1 &= \frac{-6x_1}{(1+x_1^2)^2} + 2x_2 \\ \dot{x}_2 &= \frac{-2(x_1+x_2)}{(1+x_1^2)^2}\end{aligned}$$

Proof Idea

Assume $x(t) \neq 0$ (otherwise we have $x(\tau) = 0$ for all $\tau > t$). Then

$$\frac{\dot{V}(x)}{V(x)} \leq -\alpha$$

Integrating from 0 to t gives


$$\log V(x(t)) - \log V(x(0)) \leq -\alpha t \Rightarrow V(x(t)) \leq e^{-\alpha t} V(x(0))$$

Hence, $V(x(t)) \rightarrow 0$, $t \rightarrow \infty$.

Using the properties of V it follows that $x(t) \rightarrow 0$, $t \rightarrow \infty$.

LaSalle's Invariant Set Theorem

Theorem Let $\Omega \subseteq \mathbb{R}^n$ compact invariant set for $\dot{x} = f(x)$. Let $V : \Omega \rightarrow \mathbb{R}$ be a C^1 function such that $\dot{V}(x) \leq 0$, $\forall x \in \Omega$, $E := \{x \in \Omega : \dot{V}(x) = 0\}$, $M :=$ largest invariant subset of E
 $\Rightarrow \forall x(0) \in \Omega$, $x(t)$ approaches M as $t \rightarrow +\infty$



Note that V must **not** be a positive definite function in this case.

Lyapunov Theorem for Global Asymptotic Stability

Theorem Let $\dot{x} = f(x)$ and $f(0) = 0$.

If there exists a C^1 function $V : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

- (1) $V(0) = 0$
- (2) $V(x) > 0$, for all $x \neq 0$
- (3) $\dot{V}(x) < 0$ for all $x \neq 0$
- (4) $V(x) \rightarrow \infty$ as $\|x\| \rightarrow \infty$

then $x = 0$ is globally asymptotically stable.

Somewhat Stronger Assumptions

Theorem: Let $\dot{x} = f(x)$ and $f(0) = 0$. If there exists a C^1 function $V : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

- (1) $V(0) = 0$
- (2) $V(x) > 0$ for all $x \neq 0$
- (3) $\dot{V}(x) \leq -\alpha V(x)$ for all x
- (4) $V(x) \rightarrow \infty$ as $\|x\| \rightarrow \infty$


then $x = 0$ is globally **exponentially** stable.

Invariant Sets

Definition: A set M is called **invariant** if for the system

$$\dot{x} = f(x),$$

$x(0) \in M$ implies that $x(t) \in M$ for all $t \geq 0$.



Special Case: Global Stability of Equilibrium

Theorem: Let $\dot{x} = f(x)$ and $f(0) = 0$. If there exists a C^1 function $V : \mathbb{R}^n \rightarrow \mathbb{R}$ such that

- (1) $V(0) = 0$, $V(x) > 0$ for all $x \neq 0$
- (2) $\dot{V}(x) \leq 0$ for all x
- (3) $V(x) \rightarrow \infty$ as $\|x\| \rightarrow \infty$
- (4) The only solution of $\dot{x} = f(x)$, $\dot{V}(x) = 0$ is $x(t) = 0 \forall t$

$\Rightarrow x = 0$ is globally asymptotically stable.

A Motivating Example (cont'd)

$$m\ddot{x} = -b\dot{x} - k_0x - k_1x^3$$

$$V(x) = (2m\dot{x}^2 + 2k_0x^2 + k_1x^4)/4 > 0, \quad V(0,0) = 0$$

$$\dot{V}(x) = -b|\dot{x}|^3$$

Assume that there is a trajectory with $\dot{x}(t) = 0$, $x(t) \neq 0$. Then

$$\frac{d}{dt}\dot{x}(t) = -\frac{k_0}{m}x(t) - \frac{k_1}{m}x^3(t) \neq 0,$$

which means that $\dot{x}(t)$ can not stay constant.

Hence, $\dot{V}(x) = 0 \iff x(t) \equiv 0$, and LaSalle's theorem gives global asymptotic stability.


Example—Stable Limit Cycle

$$E = \{x \in \Omega : \dot{V}(x) = 0\} = \{x : \|x\| = 1\}$$


$M = E$ is an invariant set, because

$$\frac{d}{dt}V = -2(x_1^2 + x_2^2 - 1)(x_1^2 + x_2^2) = 0 \quad \text{for } x \in M$$

We have shown that M is a asymptotically stable limit cycle (globally stable in $R - \{0\}$)



Adaptive Noise Cancellation by Lyapunov Design





$$\dot{x} + ax = bu$$

$$\dot{\hat{x}} + \hat{a}\hat{x} = \hat{b}u$$

Introduce $\tilde{x} = x - \hat{x}$, $\tilde{a} = a - \hat{a}$, $\tilde{b} = b - \hat{b}$.

Want to design adaptation law so that $\tilde{x} \rightarrow 0$

Results



Estimation of parameters starts at t=10 s.

Example—Stable Limit Cycle

Show that $M = \{x : \|x\| = 1\}$ is a asymptotically stable limit cycle for (almost globally, except for starting at $x=0$):

$$\dot{x}_1 = x_1 - x_2 - x_1(x_1^2 + x_2^2)$$

$$\dot{x}_2 = x_1 + x_2 - x_2(x_1^2 + x_2^2)$$

Let $V(x) = (x_1^2 + x_2^2 - 1)^2$.

$$\begin{aligned} \frac{dV}{dt} &= 2(x_1^2 + x_2^2 - 1) \frac{d}{dt}(x_1^2 + x_2^2 - 1) \\ &= -2(x_1^2 + x_2^2 - 1)^2(x_1^2 + x_2^2) \leq 0 \quad \text{for } x \in \Omega \end{aligned}$$

$\Omega = \{0 < \|x\| \leq R\}$ is invariant for $R = 1$.

A Motivating Example (revisited)

$$m\ddot{x} = -b\dot{x} - k_0x - k_1x^3$$

$$V(x, \dot{x}) = (2m\dot{x}^2 + 2k_0x^2 + k_1x^4)/4 > 0, \quad V(0,0) = 0$$

$$\dot{V}(x, \dot{x}) = -b|\dot{x}|^3 \text{ gives } E = \{(x, \dot{x}) : \dot{x} = 0\}.$$

Assume there exists $(\bar{x}, \dot{\bar{x}}) \in M$ such that $\bar{x}(t_0) \neq 0$. Then

$$m\ddot{\bar{x}}(t_0) = -k_0\bar{x}(t_0) - k_1\bar{x}^3(t_0) \neq 0$$

so $\dot{\bar{x}}(t_0+) \neq 0$ so the trajectory will immediately leave M . A contradiction to that M is invariant.

Hence, $M = \{(0,0)\}$ so the origin is asymptotically stable.

Let us try the Lyapunov function

$$V = \frac{1}{2}(\tilde{x}^2 + \gamma_a \tilde{a}^2 + \gamma_b \tilde{b}^2)$$

$$\begin{aligned} \dot{V} &= \tilde{x}\dot{\tilde{x}} + \gamma_a \tilde{a}\dot{\tilde{a}} + \gamma_b \tilde{b}\dot{\tilde{b}} = \\ &= \tilde{x}(-a\tilde{x} - \tilde{a}\hat{x} + \tilde{b}u) + \gamma_a \tilde{a}\dot{\tilde{a}} + \gamma_b \tilde{b}\dot{\tilde{b}} = -a\tilde{x}^2 \end{aligned}$$

where the last equality follows if we choose

$$\dot{\tilde{a}} = -\dot{\hat{a}} = \frac{1}{\gamma_a}\tilde{x}\hat{x} \quad \dot{\tilde{b}} = -\dot{\hat{b}} = -\frac{1}{\gamma_b}\tilde{x}u$$


Invariant set: $\tilde{x} = 0$.

This proves that $\tilde{x} \rightarrow 0$.

(The parameters \tilde{a} and \tilde{b} do not necessarily converge: $u \equiv 0$.)

Demonstration if time permits

Results



Estimation of parameters starts at t=10 s.

- Stability analysis using input-output (frequency) methods

<http://www.math.spbu.ru/NDA2007/en/>
International Congress "Nonlinear Dynamical Analysis - 2007". I. ...

[Russian Version](#)

International Congress "Nonlinear Dynamical Analysis - 2007"
dedicated to the 150th anniversary of Academician A.M. Lyapunov



2007 June 4-8
Russian Academy of Sciences
Saint-Petersburg State University, Russia

Second Announcement

[Preliminary registration form](#)

[Application for participation & Abstract Requirements](#)

[Registration fee](#)

[Congress Service: accommodation, visa support, transfer, cultural programme](#)

Scientific Committee of the Congress

Chair: academician **Yu.S. Osipov** - (Moscow)
Vice-chair: prof. **A.A. Farshteyn** - (Moscow), academician **V.G. Peshekhonov** - (St.-Petersburg), prof. **L.A. Verbitskaya** - (St.-Petersburg)

Members of the Scientific Committee

academician N.A. Anfinov (Moscow)	academician V.M. Matrosov (Moscow)
prof. P. Borne (France)	prof. H. Mijagi (Japan)
academician V.V. Kozlov (Moscow)	academician N.F. Morozov (St.-Petersburg)
academician N.N. Krasovskiy (Yekaterinburg)	academician V.V. Rumyantsev (Moscow)
prof. V. Lakshmikantham (USA)	prof. E.A. Tropp (St.-Petersburg)
prof. G.A. Leonov (St.-Petersburg)	academician V.Ye. Fozov (Moscow)
prof. S.N. Mamonov (Moscow)	academician K.V. Frolov (Moscow)

International Program Committee (IPC)

Co-chair: **V. Lakshmikantham** (USA), **V.M. Matrosov** (Russia)
Vice-chair: **V. A. Pliss** (Russia).

Members:

N.V. Anbedev (Russia), **A.A. Ashimov** (Kazakhstan), **V. V. Beletsky** (Russia), **P. Borne** (France), **S.N. Vasiljev**, **V.G. Veretennikov** (Russia), **V. Vojcic** (Yugoslavia), **A.V. Kamperman**, **D.M. Kimov**, **V.V. Kozlov**, **R. Kuzdrev**, **V.B. Koltunovskiy** (Russia), **V. Kretschmer**, **V.A. Kuznetsov** (Ukraine), **P.S. Krut'nikov**, **V.P. Legostayev**, **G.A. Leonov** (Russia), **S. Leela** (USA), **A.M. Ljapunov**, **A.M. Maruyama**, **I.V. Matrosov** (Russia), **H. Mijagi** (Japan), **N.F. Morozov**, **Yu.S. Osipov** (Russia), **M. Boudol** (France), **V.V. Rumyantsev**, **I.A. Rozhnov** (Russia), **A.V. Sushchik** (Russia).